
Dietary Support Guide



Omega Diagnostics defines food sensitivity as an IgG antibody reaction to food. Food IgG antibody testing is not intended to diagnose or treat any medical conditions.


Food sensitivity tests do not identify IgE-mediated food allergies or provide information about coeliac disease, enzyme deficiencies such as lactose, histamine, tyramine or alcohol intolerance, or other chemical sensitivities, such as reactions to certain food additives.

We do not recommend testing during pregnancy or breastfeeding.

Following a test, the assistance of a professional healthcare provider is advised, and any medical concerns should be referred to a medical doctor.

Omega Diagnostics is not responsible for any misinterpretation or misuse of the information that we provide, including any diagnoses or nutritional changes initiated by a healthcare practitioner or patient based upon the content of our test results. Any substantial dietary change should be conducted under the proper guidance of a qualified and registered nutritional therapist or nutritionist. This is particularly important if the individual is a child. This test is not recommended for children under 2 years of age.

This guidebook is intended for educational purposes only and the information contained within it is not intended, in any way, to be used as a diagnostic tool.



The following information is contained within this Guidebook

	Page
Introduction	5
Before changing your diet	6
How can this test help me?	7
What symptoms may improve?	8
Antibody levels	9
Test report	10
Planning and implementing my programme	12
Removing and rotating foods in your diet	14
Food replacements	15
Egg replacement	16
Wheat replacement	17
Gluten replacement	19
Yeast replacement	20
Repair and restore gut microbiome	22
Cross reactions	23
Hidden foods	24
Reintroduction of eliminated foods	25
Food reintroduction diary	26
Understanding food sensitivity	27
Immune-mediated reactions	28
Non immune-mediated reactions	29
FAQ's	30

Action plan timeline once you have your results

Interpret results	Read this guidebook
Plan the diet	Foods to remove and rotate and new foods to experiment with
Follow for 3 months	Focus on repairing the gut
Systematic food reintroduction	Reintroduce foods one at a time; one new food every 5 days
Follow new diet and retest in 9-12 months if needed	Enjoy your new varied diet

For more information seek advice from your healthcare professional.



1. Introduction



This guidebook is designed to give you advice on how to successfully change your diet based on the results of your Food Detective® test. The information contained here will help you identify which foods should be temporarily eliminated and which should be reduced or rotated and will provide ideas for replacement foods.

To understand your results and what they mean, it is important to familiarise yourself with the terminology used and what we are referring to when we discuss food sensitivity and the difference between food sensitivity and food allergy.

Food Detective® is a food sensitivity test, which identifies IgG antibodies and immune-mediated reactions to food. It is also known as a type III allergy as opposed to a type I IgE food allergy. IgG tests and IgE tests cannot be compared. IgE food allergic reactions are instant and may be life threatening, whereas IgG food sensitivity reactions may take up to 72 hours to appear and are more difficult to detect.

Please be aware that if you have a classic IgE allergy to a food this will not be reflected in your Food Detective® results, and you must always avoid that food as a priority.



Throughout this guidebook we will refer to an IgG immune-mediated reaction as a food sensitivity. As we know you are keen to get started straight away, we are outlining the most relevant points first. Reading them will help you get the most benefit from the dietary changes you make. However, if you want to delve in deeper please look out for a box like this which will show you where to head for more info on a particular topic:

[Learn more on this. See Page X](#)

2. Before changing your diet



Any substantial dietary change should be conducted under the proper guidance of a qualified and registered nutritional therapist or nutritionist. This is particularly important if the individual is a child.



If you have a medical condition, or taking medication, it is advisable to discuss your proposed dietary change with a health professional e.g. a doctor, nutritionist or dietician.

If you have previously been diagnosed with IgE food allergy, then you should continue to avoid those foods even if the food is negative on your Food Detective® test. This also applies if you are lactose intolerant and need to limit dairy consumption, or if you are sensitive to compounds found in foods such as nitrates, sulphates, or histamine.

Plan and organise your meals in advance as much as possible. By collecting recipe ideas based on your negative foods and shopping ahead you are less likely to struggle with what you eat.

Know the range of foods you can eat. While you may be sensitive to a few foods, there will be many negative foods that you should be free to eat, it is just as important to focus on the high number of foods that can be eaten as on those foods to eliminate and rotate.

When eliminating a food from your diet, simply replace with other foods from the same food group.

Recognise what food products contain your positive foods. Many ready-made meals and sauces contain a variety of ingredients that you may not have necessarily associated with the product, so it is important to always check the labels.

It is very important to maintain a healthy, nutritious diet. By eating a variety of foods, you will obtain a wide variety of nutrients and reduce the risk of further sensitivities.

3. How can this test help me?

The aim of an IgG dietary elimination programme is to help reduce symptoms that may be triggered by problematic foods and to help support the body to improve tolerance to these foods once symptoms have subsided.

This allows foods to then be reintroduced as part of a healthy, balanced diet and to be tolerated at a reduced intake.

Research has linked IgG food antibody reactions to symptoms, including irritable bowel and migraine. It may take up to 72 hours to notice symptoms, which is why it can be hard to detect culprit foods.

The production of IgG antibodies to food proteins is just one of the ways in which the body's immune system reacts to substances that adversely affect it. Once culprit foods have been identified using a food sensitivity test, adapting the diet may help to reduce any associated symptoms.

If symptoms do not reduce it is important to understand that not all food-related symptoms are caused by an IgG response; they can be caused by several other triggers, including enzyme deficiency or chemical sensitivity, by inadequate digestion, dysbiosis, candidiasis, parasites, intestinal infections, a poorly balanced diet, excess exercise, alcohol consumption, or the effects of drugs and medications. A healthcare professional will help to identify which area to focus on next.

4. What symptoms may improve?



Research shows strong evidence for improvement in Irritable Bowel Syndrome (IBS) and migraine which includes the symptoms listed below:

Symptoms associated with Irritable Bowel Syndrome include:

- ✓ Bloating
- ✓ Constipation
- ✓ Diarrhoea
- ✓ Flatulence
- ✓ Lethargy
- ✓ Nausea
- ✓ Stomach cramps / abdominal pain

Symptoms associated with migraine include:

- ✓ Headaches
- ✓ Nausea
- ✓ Vomiting
- ✓ Lethargy



Anecdotal evidence from clinical practitioners in the field suggests that food sensitivity testing may contribute to a wider variety of symptoms, but more research is needed to confirm this.

Learn more on the mechanism behind food sensitivity

See Appendix page 27




5. The Food Detective test



A food sensitivity test assesses the level of food antibodies in a blood specimen, detecting which foods are potentially provoking symptoms.

A Food Detective test assesses your IgG antibody response to 59 foods (including mixes).

The results are represented as a colour; white, pale, mid or strong. Once your practitioner has carried out your test, the wells containing the foods you have reacted to will change colour depending on the reactivity. An example of what a results tray looks like after completing the test is below:



Reaction tray diagram

Oat	Wheat	Rice	Corn	Rye	Durum Wheat
Gluten	Almond	Brazil Nut	Cashew Nut	Tea	Walnut
Milk (Cow)	Whole Egg	Chicken	Lamb	Beef	Pork
White Fish Mix ¹	Freshwater Fish Mix ²	Tuna	Shellfish Mix ³	Broccoli	Cabbage
Carrot	Leek	Potato	Celery	Cucumber	Peppers ⁴
Legume Mix ⁵	Grapefruit	Melon Mix ⁶	Peanut	Soya Bean	Cocoa Bean
Apple	Blackcurrant	Olive	Orange & Lemon	Strawberry	Tomato
Ginger	Garlic	Mushroom	Yeast	Negative control	Positive control

Mixes include:

WHITE FISH MIX: Haddock, Cod, Plaice

FRESHWATER FISH MIX: Salmon, Trout

SHELLFISH MIX: Shrimp, Prawn, Crab, Lobster, Mussel

PEPPERS: Red, Green, Yellow

LEGUME MIX: Pea, Lentil, Haricot

MELON MIX: Cantaloupe, Watermelon

If you are experiencing adverse symptoms and the Food Detective test has identified any reactive foods due to raised IgG antibody levels, this may indicate a sensitivity to those specific foods. Removing them from the diet may result in an improvement of symptoms.

Please refer to 'Planning Your Diet' for more detailed information about removal/substitution of foods.

6. Interpreting your test report



Your practitioner will explain your results to you.

WELLS 1-46

Strong colour indicates a positive reaction.

Eliminate these foods from the diet for 3 months.

Mid-colour indicates a moderate positive reaction.

Reduce these foods to once every 4 days. If there are no strong coloured reactions, eliminate foods showing a mid-coloured reaction for 3 months.

Pale colour indicates a weak positive reaction.

Observe the effect of eating these foods. If there are no other positive reactions, eliminate foods showing a pale coloured reaction for 3 months.

White colour consistent with the negative control indicates a negative reaction.

Unless the patient is allergic to these foods, they can remain in the diet as normal.

Cross-reactivity

Occasionally foods show up in a food report that have not knowingly been eaten. If the food is not hidden in the diet, then this may be due to what is known as a cross-reaction.

A cross-reaction occurs when different foods possess almost identical protein structures and an IgG specific antibody reacts to the wrong, but almost identical, protein found in food. This causes a falsely elevated result for this food. If this occurs the result should be ignored, and a healthcare professional will be able to give guidance.



To learn more on the mechanism behind cross reactions
See Appendix page 23



“

A cross reaction occurs when different foods possess almost identical protein structures

7. Planning and implementing my programme



Read the nutritional information in this section carefully before you start. We advise you to seek professional advice to ensure you maintain a balanced diet. The programme consists of 5 stages:

Step 1. REMOVE...strong positive foods for 3 months

If you have many strong positive foods, eliminate the top 4 or 5 foods which you eat most frequently. Rotate the remaining positive foods (see Step 2. ROTATE).

If no strong positives present eliminate foods showing a moderate positive reaction for 3 months.

Step 2. ROTATE

Moderate and weak positive foods for 3 months

How do you rotate foods?

- ✓ To rotate foods, eat them no more than once every 3-4 days.

Step 3. REPLACE...foods with alternative

How can you make it easy to replace foods with alternatives?

- ✓ Take a day or two to find alternative foods
- ✓ Collect recipes, make a menu plan
- ✓ Careful planning helps compliance
- ✓ Keep diet balanced and varied, for example, instead of wheat, choose oats and quinoa



Step 4. REPAIR...gut health**What does repairing the gut mean?**

Particularly important when lots of positives present. Include gut supportive supplementation and functional foods:

- ✓ Probiotics
- ✓ Bone broth
- ✓ Fermented foods, such as sauerkraut
- ✓ Oily fish, such as salmon
- ✓ Root vegetables, such as sweet potatoes
- ✓ Inulin rich vegetables, such as leeks and bananas

Step 5. REINTRODUCTION...of

foods eliminated after 3 months

Do the foods need to be avoided forever?

- ✓ No. Reintroduce foods gradually - 1 portion every 5 days x 1 food at a time
- ✓ Start with the weak positive foods
- ✓ Monitor symptoms before reintroducing the next food
- ✓ Avoid for longer if necessary
- ✓ Eat foods in moderation and continue to vary the diet
- ✓ Retest if needed after 1 year - don't re-test too soon

If symptoms have not improved after 3 months despite strictly following results in your Food Detective® test, this could indicate that IgG-mediated food sensitivity is not the only cause of your symptoms.

We recommend that you consult with your health practitioner to investigate other potential causes. This could include, amongst others, imbalanced microbiome, lifestyle triggers, e.g., stress and the impact it has on the gut and enzyme deficiencies.

To learn more see Appendix page 22



8. Removing and rotating foods in your diet



If you have reacted to more than 10 foods, we recommended to avoid the 4-5 which you eat most frequently. The remaining positive foods may be eaten in moderation, e.g., every 3-4 days.

In order to prevent new sensitivities developing, do not replace any positive foods exclusively with a single alternative.

Focus on variety when selecting alternative foods. For example, if wheat is being eliminated from the diet, try a different breakfast every day rather than simply swapping toast for porridge oats, or alternate between a few different options throughout the week, e.g., porridge oats one day, fruit and yoghurt the next day, and poached egg on rye bread the next day.

Rotate Foods



To rotate foods, eat them no more than once every 3-4 days. For example, to rotate wheat and dairy see image below.



Monday	Tuesday	Wednesday	Thursday	Friday	Saturday	Sunday
Wheat bread	Oat cakes	Rye bread	Wheat bread	Oat cakes	Rye bread	Quinoa crispbread
Cow's milk	Coconut milk	Rice milk	Cow's milk	Almond milk	Coconut milk	Hemp milk

In summary:

- Avoid eating any one food too regularly.
- Limit consumption of each food to no more than twice per week.
- Include a wide variety of foods in the diet to ensure that a range of important vitamins and minerals are consumed. Studies show that consumption of more than 30 different plant-based foods per week has been found to deliver health benefits and a more diverse gut microbiome.



9. How to remove: Dairy



Aim to focus on eating a wholefood diet and cooking from scratch with simple ingredients where possible. Please explore the “free from” sections in supermarkets and your local health food shops as there is an ever-increasing variety of choices.

Dairy Replacement



If the Food Detective® test has shown a positive reaction to milk, it is recommended that you eliminate all consumption of milk and milk products for three months. See below for replacement ideas and focus on your negative food list.

- ✓ Plant-based milks, e.g., coconut milk
- ✓ Dairy-free butter, e.g., vegan block butter, coconut oil
- ✓ Spreads, e.g., hummus, almond butter
- ✓ Cheese, e.g., nut cheeses, coconut cheese
- ✓ Yoghurt, e.g., coconut and oat yoghurt
- ✓ Mayonnaise, e.g., dairy-free mayonnaise
- ✓ Cream, e.g., oat cream
- ✓ Ice cream, e.g., dairy-free ice cream
- ✓ Chocolate, e.g., dark chocolate, vegan bars

Dairy-free alternatives are often highly processed, so be sure to check ingredients carefully and for other moderate or weak positive foods on your report and eat wholefoods where possible.

When removing dairy from your diet be aware that it may be hidden in some foods and so it is important to always read the food ingredient labels carefully before purchasing. Check for hidden and other names for dairy, including:

- ✓ Butter
- ✓ Casein, caseinate
- ✓ Beta-lactoglobulin
- ✓ Alpha-lactalbumin
- ✓ Milk powder, skimmed milk powder
- ✓ Milk solids, non-fat milk solids
- ✓ Whey

Replacing foods of equivalent nutritional value is vital to health. Seek nutritional advice and ensure nutrients found in dairy are replaced with:

- ✓ Calcium: seeds including sesame, chia, flax and pumpkin, canned sardines and salmon, beans and lentils, almonds, leafy green vegetables, tofu, edamame and figs.
- ✓ Protein: fish, nuts, seeds, eggs, meat, lentils and beans.
- ✓ Vitamin D: natural sunlight (expose skin to UV sunlight for 10-30 minutes daily but do not burn), mushrooms, cod liver oil or nutritional supplements.
- ✓ Vitamin A: leafy green vegetables and orange and yellow vegetables, e.g., sweet potatoes and carrots, oily fish, liver and tomatoes.



10. How to remove: Eggs



If the Food Detective® test has shown a positive reaction to eggs, it is recommended that you eliminate these from your diet for three months. See below for replacement ideas and focus on your negative list:

Egg Replacement



- ✓ Egg-free pasta, e.g., rice or buckwheat
- ✓ Egg-free noodles, e.g., rice or buckwheat
- ✓ Mayonnaise, e.g., egg-free mayonnaise
- ✓ Ice cream, e.g., sorbet or egg-free ice cream
- ✓ Replace eggs in cakes and cooking with either 1 tbsp ground flaxseed plus 3 tbsp water and allow to soak for 15 mins, or with mashed banana
- ✓ Replace egg as thickener with aquafaba (water in which chickpeas and other pulses have been cooked)
- ✓ Puddings, e.g., fresh fruit and fruit crumbles, sorbet, jelly

Egg-free alternatives are often highly processed, so be sure to check ingredients for other moderate or weak positive foods on your report and eat wholefoods where possible.

When removing eggs from your diet be aware that they may be hidden in some foods and so it is important to always read the food ingredient labels carefully before purchasing. Check for hidden and other names for eggs, including:

- | | |
|-------------------|---------------|
| ✓ Albumin | ✓ Ovalbumin |
| ✓ Egg white | ✓ Ovovitellin |
| ✓ Egg yolk | ✓ Ovaglobulin |
| ✓ Frozen egg | ✓ Ovamucin |
| ✓ Pasteurised egg | ✓ Globulin |
| ✓ Dried egg | ✓ Livetin |
| ✓ Egg powder | ✓ Vitellin |
| ✓ Egg protein | |

Replacing foods of equivalent nutritional value is vital to health. Seek nutritional advice and ensure nutrients found in eggs are replaced with:

- Protein: fish, nuts, seeds, eggs, meat, lentils and beans.
- Vitamin D: natural sunlight (expose skin to UV sunlight daily for 10-30 minutes, but do not burn), mushrooms, cod liver oil or supplements.
- Vitamin A: leafy green vegetables and orange and yellow vegetables, e.g., sweet potatoes and carrots, oily fish, liver and tomatoes.
- Selenium: Brazil nuts, fish, meat, sunflower seeds, mushrooms.
- B2: beef, tofu, milk, fish, mushrooms, pork, spinach, almonds, avocados.



11. How to remove: Wheat



If the Food Detective® test has shown a positive reaction to wheat, it is recommended that you eliminate consumption of wheat and wheat products for three months. See below for replacement ideas and focus on your negative list.

- ✓ Bread, e.g., rice bread, corn bread, 100% rye bread
- ✓ Pasta and noodles, e.g., buckwheat or rice
- ✓ Cereals, e.g., oat muesli, sugar-free cornflakes
- ✓ Sauces and gravies, e.g., arrowroot, cornflour
- ✓ Flours, e.g., buckwheat, almond flour
- ✓ Biscuits, e.g., maize or oat based
- ✓ Processed meat, e.g., wheat-/gluten-free alternatives
- ✓ Soy sauce, e.g., coconut aminos, tamari

Wheat-free alternatives are often highly processed, so be sure to check ingredients for other moderate or weak positive foods on your report and eat wholefoods where possible.

Replacing foods of equivalent nutritional value is vital to health. Seek nutritional advice and ensure nutrients found in wheat are replaced with:

- Fibre: non-processed foods, such as brown rice, quinoa, buckwheat, oats, root vegetables with skins washed and not peeled, plenty of fruits and vegetables daily.
- Starch: maize, millet, rice and quinoa, lentils, beans, potatoes, sweet potatoes, parsnips, plantains, green bananas and yams.
- Protein: nuts, seeds, eggs, meat, fish and eggs.
- B vitamins (particularly relevant for a meat-free diet): black-eyed beans, brown rice, Marmite, salmon, tofu, avocado, spinach and peas.



When removing wheat from your diet be aware that it may be hidden in some foods and so it is important to always read the food ingredient labels carefully before purchasing. Check for hidden and other names for wheat, including:

- | | | |
|--------------------------|--------------------------------|---------------------------|
| ✓ Abyssinian hard | ✓ Farro | ✓ Matzo/Matzoh/Matzah |
| ✓ Atta | ✓ Filler | ✓ Modified food starch |
| ✓ Barley | ✓ Flour | ✓ Modified starch |
| ✓ Binder or binding | ✓ Food starch | ✓ Rusk |
| ✓ Bran | ✓ Frumento | ✓ Rye |
| ✓ Bulgur | ✓ Fu | ✓ Seitan |
| ✓ Cake flour | ✓ Graham flour | ✓ Semolina |
| ✓ Cereal | ✓ Groats | ✓ Sooji/Suji |
| ✓ Cereal binders/binding | ✓ Gum base/dextrin | ✓ Special edible starch |
| ✓ Cereal protein | ✓ Hydrolysed wheat protein | ✓ Spelt |
| ✓ Couscous | ✓ Hydrolysed vegetable protein | ✓ Starch |
| ✓ Dinkel | ✓ Kamut | ✓ Thickener or thickening |
| ✓ Durum | ✓ Maida | ✓ Triticale |
| ✓ Edible starch | ✓ Malt | ✓ Triticum |
| ✓ Einkhorn | ✓ Malt extract (wheat-based) | ✓ Wheat |
| ✓ Emmer | ✓ Manna | ✓ Wheat bran hydrolysate |
| ✓ Enriched flour | | ✓ Wheat protein isolate |
| ✓ Farina | | |

REMEMBER: always check the label. Please note that products labelled gluten-free may not be wheat-free as some are made from wheat starch and these are not suitable for wheat-free diets.



12. How to remove: Gluten



If the Food Detective® test has shown a positive reaction to gluten it is recommended that you eliminate consumption of all grains containing gluten; these include wheat, see above for wheat varieties, for three months. We isolate gluten separately from the other proteins in grains so if gluten is positive all gluten containing grains should be avoided even if those individual grains, e.g., rye, show as negative on your report. See below for replacement ideas.

Gluten-free alternatives are often highly processed, so be sure to check ingredients for other moderate or weak positive foods on your report and eat wholefoods where possible.

When removing gluten from your diet be aware that it is contained in all the wheat varieties listed on the previous page so these foods should be avoided. It is important to always read the food ingredient labels carefully before purchasing.

Gluten Replacement



- ✓ Naturally gluten-free flours: amaranth, potato, quinoa, buckwheat, rice, corn, quinoa, ground nuts, e.g., almonds and pecan, lentil flour, chickpea/gram flour, soy flour, millet, tapioca
- ✓ Gluten-free bread, e.g., rice bread or corn bread
- ✓ Gluten-free pasta and noodles, e.g., buckwheat or rice
- ✓ Gluten-free cereals, e.g., quinoa porridge, sugar-free cornflakes
- ✓ Sauces and gravies, e.g., arrowroot, cornflour
- ✓ Flours, e.g., chickpea, lentil
- ✓ Biscuits, e.g., maize or gluten-free oat cakes
- ✓ Processed meat, e.g., gluten-free alternatives
- ✓ Soy sauce, e.g., coconut aminos, tamari
- ✓ Soy sauce, e.g., coconut aminos, tamari

Please note that oats are gluten-free but are often contaminated with wheat, rye and barley during processing. It is therefore recommended that foods containing oats are also avoided unless they are labelled gluten-free.



13. How to remove: Yeast



If the Food Detective® test has shown a positive reaction to yeast, it is recommended that you eliminate consumption of both baker's and brewer's yeast foods for three months. See below for replacements.



Of all the foods to avoid, baker's and brewer's yeast, being two strains of the same organism *Saccharomyces cerevisiae*, are probably the most difficult as they are used in the manufacture of such a wide variety of foods due to the yeast's excellent food safety record.

It is used in most bread-making, brewing and fermentation processes. Baker's yeast is chosen for its flavour and ability to make carbon dioxide, while brewer's yeast has a more bitter taste and lends itself to producing alcohol.

Yeast replacement

- ✓ Yeast-free bread, e.g., flatbreads, matzos, soda bread, pumpernickel bread
- ✓ Yeast-free biscuits, e.g., rice cakes, tortillas, oat biscuits
- ✓ Baking, e.g., baking powder, bicarbonate of soda
- ✓ Sauces and gravies, e.g., arrowroot, yeast-free stock cubes, tamari
- ✓ Yeast-free alcohol, e.g., spirits and prosecco
- ✓ Fermented foods, e.g., sauerkraut, tempeh, kimchi



Yeast-free alternatives are often highly processed, so be sure to check ingredients for other moderate or weak positive foods on your report and eat wholefoods where possible.

When removing yeast from your diet be aware that it is contained as an ingredient in many foods, and so it is important to always read the food ingredient labels carefully before purchasing. Below is a list of foods and ingredients to look out for:

- ✓ Baker's yeast in leavened baked goods, e.g., breads, muffins, croissants, biscuits, and other baked goods
- ✓ Yeast extract, e.g., Marmite, Vegemite, Bovril, stock cubes and gravies
- ✓ Fermented food and drink, e.g., beer, wine, cider, ginger ale, soy sauce, kombucha, kefir, miso, dressings
- ✓ Malt syrup, malt extract

Replacing foods of equivalent nutritional value is vital to health. Seek nutritional advice and ensure nutrients found in yeast are replaced with:

- ✓ B vitamins: meat, fish, whole grains, nuts, seeds and dark green leafy vegetables.
- ✓ Selenium: Brazil nuts, tinned fish and shellfish, beef, turkey, chicken, fortified cereals, wholewheat bread, beans and lentils.
- ✓ Chromium: beef and animal proteins, wholewheat flour, grape juice, tomato juice, apples and green beans.



IMPORTANT: We are testing for the protein in yeast, we are not testing for Candida albicans. You should therefore only avoid foods containing baker's and brewer's yeast. This is not an anti-candida diet, and the results of this test cannot tell you if you have a candida overgrowth.

14. Repair gut wall barrier and restore gut microbiome



Testing positive to a high number of foods could indicate increased intestinal permeability or “leaky gut”. This is a condition used to describe a loss in function of the epithelial (gut) wall, which is highly selective in terms of what it allows to be absorbed into the body.

Several factors can contribute to development of ‘leaky gut’. These can include lifestyle, poor diet, medications, and stress, resulting in damage and dysbiosis due to a reduction in microbial diversity, including the loss of beneficial bacteria. These can all increase permeability of the gut wall allowing food particles to cross into the bloodstream before they are adequately digested, provoking an IgG immune response.

The resulting response triggers the production of complement which in turn triggers inflammation and symptoms.

While following an IgG elimination diet, with the resulting reduction of IgG immune responses and complexes, it is also important to take steps to repair the gut wall so that positive foods can be reintroduced without provoking a reaction.

The dietary recommendations below may help to improve “leaky gut” and any associated inflammation, allowing the gut to heal and setting the stage for the reintroduction phase of the programme. Please seek support from an expert when navigating this stage.

Eat



- ✓ A diet rich in nutrients including vitamins A, C, E and zinc, such as the Mediterranean-style diet, which includes fish, fruit, vegetables, nuts, seeds wholegrains and olive oil.
- ✓ Homemade bone broth which is rich in gut-healing nutrients.
- ✓ Foods high in anti-inflammatory Omega 3 fats, such as salmon, sardines, walnuts, flaxseed and pecans.
- ✓ A quality probiotic supplement.
- ✓ Probiotic foods, such as pickles, sauerkraut, miso soup and kimchi.
- ✓ Many other supplements have been found to help repair the gut barrier wall. Speak to your healthcare practitioner for more information on the most beneficial healing supplements tailored to your needs.

Reduce



- ✓ Eliminate strong positive foods.
- ✓ Rotate moderate and weak positive foods.
- ✓ Reduce foods high in saturated fat, e.g., red meat and butter.
- ✓ Reduce sugar intake found in biscuits, sweets, and cakes.
- ✓ Restrict processed foods and fast food high in proinflammatory fats and antinutrients.
- ✓ Limit caffeine as it is a gut irritant.
- ✓ Reduce high intake of alcohol and excessive exercise.

15. Cross-reactions



Cross-reactivity occurs when an antibody recognises not only the food protein, or antigen, for which it was originally formed, but also others that have a significant degree of similarity to that of the original antigen. Cross-reactions are the degree to which various antigens appear alike to the immune system.

This structural similarity enables an antibody, specifically raised against one particular antigen, to recognise another antigen as being identical and to bind with it. An individual who has elevated IgG antibodies to a specific food may therefore show raised antibodies to other foods with similar protein antigens in an IgG-mediated food sensitivity test. It is not uncommon for this to occur, and it is important to understand as it can be confusing if a food is shown as positive but is not included within the diet.

While cross-reactions within the same food family may occur, it must not be assumed that this will be the case. Therefore, only foods shown as positive in the test results should be avoided and any other potentially cross-reactive foods should be kept in the diet. It is also possible that antibodies produced against a specific food may cross-react with non-food items, such as dust mites, pollen, latex or animal dander. This is known as concomitant food sensitivity. What this means is if a food tests positive in a Food Detective® test, but is not included in the diet, it should continue to be avoided whilst the other foods are temporarily removed from the diet.



Antibodies produced against a specific food may cross-react with non-food items, such as dust mites, pollen, latex or animal dander.



16. Hidden Foods



Sometimes foods are eaten unknowingly, and the Food Detective® test can throw up a positive reaction to a food ingredient which may come as a surprise.

Some examples are within cosmetics, wherein products, particularly lipsticks, often contain animal proteins, and can sometimes be the cause of a positive reaction to a specific animal protein.

If processed foods are eaten regularly, hidden foods may be consumed unknowingly. The simplest way of avoiding hidden food ingredients is to eat homemade meals so that the ingredients are known.



17. Reintroduction of Eliminated Foods



After at least 3 months, and only when symptoms have subsided, positive foods may be reintroduced to the diet. However, this should be a gradual process and planning is essential.

Start with the weak positive foods and eat one portion per day for 5 days. Introduce one food at a time and monitor your symptoms over the 5-day period. If symptoms return, this food is still likely to be a problem and you should revert to the elimination diet for a further two months before a second reintroduction challenge.

Reintroduce the weak positive foods first (one portion per day for 5 days)



Leave 5 days before attempting the next food reintroduction



No Reaction

This food can now be added back into the diet



Symptoms Experienced

Continue IgG elimination diet for another 1-2 months, focusing on gut health before attempting reintroduction

If symptoms do not return when a food is reintroduced, this food can be included in your diet, but eaten only occasionally in small amounts. A second food can then be reintroduced and symptoms monitored for 5 days, and so on until you have tested all your positive foods.

Example Food Reintroduction Diary:

Reintroduced food	Date of first reintroduction	Symptom	Date of second reintroduction
Orange	01/07/2020	None	N/A
Oat	06/07/2020	None	N/A
Cow's Milk	11/07/2020	Diarrhoea, bloating, nausea	11/09/2020

19. Understanding Food Sensitivity



Terminology

The terms 'food allergy', 'food intolerance' and 'food sensitivity/hypersensitivity' are often used interchangeably and this can be quite confusing. In simple terms, they all mean that an individual has reacted to a food that has been consumed and this then manifests as symptoms. The reactions they trigger can also vary greatly. This includes the symptoms experienced and how quickly the symptoms manifest after the food has been consumed, as well as the severity of the symptoms and how long they last, and if they are life threatening.

Some of these reactions are immune-mediated and involve the immune system, including the production of IgE and IgG antibodies. Other reactions do not involve the immune system and are non-immune mediated.



What is the immune system?

To help us understand these varying reactions, it can be helpful to have some basic knowledge of the immune system and how it works. The immune system is made up of various organs, cells and proteins and its role is to protect the body from harmful substances, germs, pathogens, fungi, and cell alterations that can result in illness. Essentially, the main functions of the immune system are:

- ✓ To fight pathogens, e.g., viruses, bacteria, or parasites and remove them from the body.
- ✓ To identify and neutralise harmful substances from the environment.
- ✓ To fight disease-causing alterations in the body, e.g., cancer cells

The immune system can be activated by substances that it doesn't recognise as being a part of the body. Common examples include proteins found on the surfaces of microbes, such as bacteria, fungi, and viruses and these are referred to as antigens. Receptors on the immune cells attach to these antigens and this then triggers an immune reaction, which includes the production of antibodies. Antibodies can then help to neutralise the microbes so they can't multiply and cause illness.

20. Immune-Mediated Reactions



Reactions that trigger an immune response are most often referred to as 'allergies' and occur when the body over-reacts to foods that do not usually produce a response in the majority of people. This overreaction triggers the immune system to produce antibodies to attack and neutralise the 'foreign' food proteins which the immune system incorrectly recognises as a threat.



Allergies are grouped into four types: I, II, III and IV. These classifications are based on which part of the immune system is activated and how long it takes for a reaction to occur. The two types of allergy that are most often associated with adverse reactions to food are:

Type I Allergy

Also known as IgE-mediated allergy/type I hypersensitivities/true allergy

These reactions are characterised by the production of IgE antibodies and the release of histamine and other chemical mediators, upon exposure to an allergen (e.g., peanuts and shellfish). They are responsible for the 'immediate-onset' of symptoms that can occur within seconds or minutes following ingestion of certain foods. Symptoms often associated with a classic 'allergic response' includes rashes, sneezing, difficulty breathing, and anaphylactic shock. It is usually obvious which foods are responsible for a food allergy and these have to be avoided for life.

Type III Allergy

Also known as IgG-mediated allergy/food-hypersensitivity

These reactions are characterised by the production of IgG antibodies and the gradual formation of antigen/antibody complexes which are deposited in tissues, leading to an inflammatory response. This type of immune reaction results in 'delayed-onset' of symptoms which can occur several hours or days after foods are ingested. It is possible to eliminate the offending food(s) from the diet for a short periods of time and then gradually reintroduce them when symptoms have improved.

21. Non Immune-Mediated Reactions



Reactions that do not produce an immune response are often referred to as 'food intolerances'. They can be caused by reactions to certain chemicals/additives found in food or, more commonly, due to enzyme deficiencies:

Enzyme Insufficiency/Deficiency

- ✓ Lactose Intolerance
- ✓ Caused by a deficiency of lactase - an enzyme that breaks down lactose (a complex sugar).
- ✓ Foods that contain lactose include: dairy products (milk, cheese, yoghurts, etc).
- ✓ Symptoms include bloating, diarrhoea and flatulence.

Histamine Intolerance

- ✓ Caused by an elevated histamine level due to a deficiency or inhibition of diamine oxidase (DAO) - an enzyme that breaks down histamine (a chemical that triggers an inflammatory response).
- ✓ Aggravated by foods high in histamine, including red wine, cheese, and tuna.
- ✓ Some foods are low in histamine but can trigger the release of histamine in the body, including citrus foods, bananas, tomatoes and chocolate.
- ✓ Symptoms include migraines, dizziness, bowel/stomach problems, rhinitis, depression, irritation and reddening of the skin.

Understanding the different reactions to food:

	IgE Type 1 Allergy	IgG-mediated food sensitivity	Non immune-mediated, e.g., enzyme insufficiency
Involves the immune system	Yes	Yes	No
Common symptoms	Hives, itchy eyes, swelling of lips/tongue, shortness of breath, nausea, vomiting, anaphylaxis	Migraines/headaches, IBS (diarrhoea, constipation, bloating, flatulence, stomach cramps, lethargy)	Gastrointestinal symptoms, hives, itchy eyes, facial swelling (dependent on specific enzyme deficiency)
Speed of reaction	Immediate (within 2 hours)	Usually 24-72 hours	Within 30 minutes to 2 hours
Severity of symptoms	Can be potentially life threatening	Can be potentially debilitating and chronic	Mild to severe
Mechanism of action	IgE antibody stimulation of histamine production	IgG antibody/antigen complex formation and complement activation	Enzyme deficiency / insufficiency
Lifelong condition	Yes	No	Yes

The Food Detective® IgG antibody test measures immune-mediated, type III allergy reactions or food sensitivities.

Frequently Asked Questions



Is it possible to be affected by foods that are not detected by the Food Detective® test?

Some foods may cause a classic allergic reaction involving the production of IgE antibodies (type I allergy) to food antigens or proteins in food. These will not be detected by the Food Detective® test as it detects IgG antibodies.

There are many foods that can cause a reaction in the body without involving the immune system, these are known as food intolerances and can produce symptoms similar to IgG reactions: amines found in chocolate, cheese and red wine may cause migraines; some food additives, such as tartrazine, can trigger hives, rashes, and asthma; monosodium glutamate (MSG) found in restaurant/take-away food can provoke sweating and dizziness; 'Nightshade' alkaloids in potatoes, tomatoes and peppers may affect the joints.

Food intolerance may be due to a deficiency of a particular enzyme, such as lactase in lactose intolerance.

Avoid foods if you suspect they are causing adverse effects.

I have been avoiding a food for several months/years. Will this affect my test results?

The Food Detective® IgG antibody test is based on the immune system's ability to produce antibodies in response to certain foods. If a food has been avoided for more than 3 months, it is possible that IgG antibody levels will be insufficient to be detected by the test and may give a negative result. To test sensitivity to a certain food, it should be included in the daily diet for five days before testing, or at least every other day, for 4-6 weeks before testing. However, if the food concerned is known to cause extreme symptoms/discomfort or you have a diagnosed allergy to it, do not reintroduce it.

If cow's milk is positive, does this mean that I am lactose intolerant?

No. Lactose intolerance is the inability to digest lactose, the major sugar found in milk, and is caused by a deficiency in the enzyme lactase. The Food Detective® test detects an IgG-mediated food sensitivity caused by the specific proteins found in milk, but does not detect the lactase enzyme and, therefore, cannot diagnose lactose intolerance.



Why do you not test for sugars?

The Food Detective® test detects level of food proteins in the blood. Sugars contain no proteins to which the IgG antibodies can combine.

Is the Food Detective® test suitable for testing children?

Yes, but we recommend a minimum age limit of 2 years to allow for the child's immune system to mature.

If my test shows reactivity towards wheat, does that mean I have coeliac disease?

No, coeliac disease is an immune disease that causes a severe reaction to the protein gluten, which is just one of the proteins found in wheat, barley, and rye. We remove gluten from our wheat extract so if wheat is positive this means you are reacting to the other wheat proteins found in this grain, and not gluten. Coeliac disease cannot be determined in a Food Detective® test, it requires a different test, should be done under medical supervision and will involve a confirmation biopsy and a positive transglutaminase response.

Is gluten-free the same as wheat-free?

No. A product can be wheat-free but not gluten-free, and vice versa. Products are available that are both gluten-free and wheat-free. It is important to read the ingredients label to be certain. The Food Detective® test uses wheat and rye food extracts that do not contain gluten. Gluten is extracted from the grains and tested separately.

If your test results show a positive reaction to gluten, it is important to eliminate any foods that contain the gluten-based grains wheat, rye and barley and substitute with naturally gluten-free foods, such as quinoa, buckwheat, corn, oats, and wild rice. If your test results show a positive result for wheat, or rye, but NOT for gluten, the reaction may be due to one of the other proteins found in the grains.

Do I need to be cautious when removing a food group from my diet?

Yes, you should be careful when introducing a new dietary regime and removing foods. We recommend seeking professional advice from registered qualified nutritionists or a dietician to help support you make these changes to ensure that nutrients are adequately replaced.

Do I need to have a re-test after a few months?

Most people do not need to re-test if symptoms have resolved, but if you would like to take another test we advise a period of 9-12 months between tests.

What if I don't experience any improvement at all?

If, after changing your diet according to the test results, improvements have not been achieved after 3 months, food sensitivity may not be the only cause of your symptoms and other investigations should be undertaken by your healthcare practitioner.



Why do we not test for IgA?

IgA is the principal isotype in secretions (especially mucus epithelium of the intestinal and respiratory tracts). It is a neutralising “first line of defence” antibody against bacteria and toxins, binding foreign antigens into complexes which are then removed by macrophages, but with little or no resultant inflammation. IgA antibodies are characterised by less specific antigen binding sites than IgG antibodies, resulting in higher cross reactivity and false positive potentials.

Why do we test for total IgG and not just IgG4?

IgG4 is an anti-inflammatory antibody triggered by IL-10. IgG4 antibodies are involved in the desensitisation of type I allergies (IgE). The guideline of the EAACI (European Academy of Allergy and Immunology) states that testing for IgG4 is not recommended for the detection of delayed food allergies (www.ncbi.nlm.nih.gov/pubmed/18489614) as it is biochemically difficult to measure. IgG4 indicates tolerance, not hypersensitivity, as it is not able to induce inflammation.

Why do we not test for anti-C3 antibody/complement?

The complement cascade is an important part of the immune system that enhances the ability of antibodies and phagocytic cells to clear microbes and damaged cells from an organism. This is, in part, driven by an inflammatory response and is beneficial to the body when fighting pathogens. Although the complement system is a component of the innate immune system, which is non-specific, it can also be recruited and brought into action by IgG antibodies generated by the adaptive immune system, which make it specific.

Some companies claim that incorporating an anti-C3d antibody improves the reliability of the food sensitivity test by establishing which foods are triggering complement and, therefore, inflammation. However, the main issue with this hypothesis is this reaction cannot take place in-vitro. Apart from technical validity of the concept issues, when writing, there were no independent published studies using this methodology.



Speak to your healthcare professional today



572-39 VERSION 1 OCTOBER 2022 ©2022 Omega Diagnostics Ltd

Omega Diagnostics defines food sensitivity as an IgG antibody reaction to food. Food IgG antibody testing is not intended to diagnose or treat any medical conditions. The Food Detective® test does not identify IgE-mediated food allergies or provide information about coeliac disease, enzyme deficiencies such as lactose, histamine, tyramine or alcohol intolerance or other chemical sensitivities such as reactions to certain food additives.

The assistance of a professional healthcare provider is advised and any medical concerns should be referred to a medical doctor. Specifications, terms and pricing are subject to change at any time. Not all products are available in all countries. Food Detective is a registered trademark of Omega Diagnostics Ltd in the UK and other countries.

References to published literature available on request | Image credit: Shutterstock

Food Detective has been developed and is manufactured by
Cambridge Nutritional Sciences Ltd, Eden Research Park, Henry Crabb Road, Littleport,
Cambridgeshire, CB6 1SE, United Kingdom